

P R A C E O R Y G I N A L N E

Nowowiejska skala nasilenia objawów alkoholowego zespołu abstynencyjnego (skala NOWA)

Nowowiejska alcohol withdrawal assessment scale (NOWA Scale)

Dariusz Wasilewski^{1,2}, Marcin Wojnar¹, Iwona Grobel²,
Grażyna Wieczorkowska³, Izabela Żmigrodzka¹, Sylwia Fudalej¹

¹Katedra i Klinika Psychiatryczna Akademii Medycznej, Warszawa

²Centrum Psychoprofilaktyki i Terapii, Warszawa

³Wydział Psychologii, Uniwersytet Warszawski

Abstract – Background. A reliable assessment of severity of alcohol withdrawal is essential for monitoring efficacy of treatment applied. Among many instruments, the Clinical Institute Withdrawal Assessment-Alcohol Scale (CIWA-A) created by Shaw et al. is most widely used. Methods. We modified and revised the CIWA-A Scale.

Results. On the basis of clinical evaluation and statistical analyses we shortened the CIWA-A to seven items, which are the most clinically pertinent. We added the item of heart rate, which is strongly related with autonomic hyperactivity.

Conclusions. NOWA Scale appears to be a valid and reliable instrument, which takes only five minutes to administer. The new scale is useful in monitoring clinical course and treatment of alcohol withdrawal.

Key words: alcohol withdrawal, delirium tremens, CIWA-A Scale

Badania finansowane przez Akademię Medyczną w Warszawie: granty na pracę własną nr IIE-18, nr IIE-67 i nr P-71.